

Maximum Run Length In a Toroidal Grid Graph

Margaret I. Doig
University of Notre Dame
mdoig@nd.edu*

February 1, 2008

Abstract

A *grid graph* is a Cartesian product $G = \gamma_1 \times \gamma_2 \times \cdots \times \gamma_k$ where the γ_i are cycles or paths. The *run length* of a Hamiltonian cycle in a grid graph is defined to be the maximum number r such that any r consecutive edges include no more than one edge in any dimension. In this paper, we present several methods for producing cycles of high run length from cycles of lower run length; sample applications include showing that the maximum run length of G is (i) less than k if each γ_i is directed; (ii) at least $\lfloor k/3 \rfloor + 1$ if each $|\gamma_i|$ has the form p^{r_i} for some fixed prime p ; and (iii) at least $\lfloor k/2 \rfloor + 1$ if every $|\gamma_i|$ is of the form $2^{q_i} p^{r_i}$ for fixed prime p .

1 Introduction

We investigate the run length of a cycle or path H which occurs in a grid graph G . A *grid graph* is the Cartesian product $G = \gamma_1 \times \gamma_2 \times \cdots \times \gamma_k$ where the γ_i are either cycles or paths. The grid graph $\gamma_1 \times \gamma_2 \times \cdots \times \gamma_k$ is said to be *k-dimensional*, and we sometimes refer to γ_i as a *dimension* of G . The *size* of the dimension is the number of vertices in γ_i . A *torus* or *toroidal grid graph* is a grid graph where each γ_i is a cycle. A grid graph is *directed* if each γ_i is directed and *undirected* if no γ_i is directed.

The *run length* of a cycle or path H in a grid graph, denoted $rl(H)$, is the maximum number r such that any sequence of r consecutive edges in H contains no more than one edge in any one dimension. The *maximum run length* of a grid graph or $mrl(G)$ is the maximum $rl(H)$ for all Hamiltonian cycles H in G , that is, for all cycles which pass through each vertex exactly once.

Run length first arose in computing applications. Since every Hamiltonian path in a torus corresponds to a listing of the coordinates of consecutive vertices, a path on a k -cube corresponds to a Gray code, i.e., a listing of all k -bit binary words in such a way that any two consecutive words differ in only one bit position. The *gap* of a Gray code is the run length of the corresponding Hamiltonian path. Electronic position-to-digital converters such as photon detectors use a Hamiltonian path to match each received

*Please send all paper correspondence to: Margaret Doig, 137 Lewis Hall, Notre Dame, IN 46556-5616




Figure 1: Composing cycles. A Hamiltonian cycle H_1 in $\gamma_1 \times \gamma_2$ is combined with H_2 from $\gamma_3 \times \gamma_4 \times \gamma_5$ to create a cycle H in $\gamma_1 \times \gamma_2 \times \cdots \times \gamma_5$. One edge is taken from the direction of H_1 , then three edges from H_2 , and so on; since both H_1 and H_2 have run length two, H has also run length two.

photon to a binary code word representing position, and machine error decreases as the code's gap increases. (See Goddyn et al. [3] for a more detailed explanation.)

The run length of Hamiltonian cycles on k -cubes was addressed in 1988 by Goddyn, Lawrence, and Nemeth [3], who investigated the gap of k -bit Gray codes and gave methods for producing them which achieve a gap that approaches $2k/3$ for large k . In 2001, Goddyn and Gvozdjak [2] placed a lower bound on the gap of a Gray code which is also a lower bound on the maximum run length of a k -cube Q_k . They showed that $mrl(Q_k)/k \rightarrow 1$ as $k \rightarrow \infty$. In addition, Ruskey and Sawada [4] thoroughly investigated the case of *bent* Hamiltonian paths and cycles, those whose run length is at least 2. If $G = \gamma_1 \times \cdots \times \gamma_k$ is a k -dimensional torus, then $mrl(G) \geq 2$ either when $k \geq 3$ or when $k = 2$ and the sizes of the dimensions are even. If G is a k -dimensional non-toroidal grid graph and $k \geq 3$, then $mrl(G) \geq 2$ if and only if the size of some dimension is even. Ruskey and Sawada also give conditions for the existence of bent Hamiltonian paths in graphs without bent cycles.

In this paper, we give lower bounds for $rl(H)$ and $mrl(G)$, where H is a Hamiltonian cycle in a torus G , by presenting several methods for producing cycles of high run length. As explained in Section 2, one method considers the Cartesian product G of tori G_1 and G_2 . A Hamiltonian cycle H can be constructed in G from Hamiltonian cycles H_1 in G_1 and H_2 in G_2 where $rl(H)$ is greater than either $rl(H_1)$ or $rl(H_2)$, and we give conditions for producing such an H with run length greater than 2. These methods may be easily applied inductively to the cases where $G = \gamma_1 \times \gamma_2 \times \cdots \times \gamma_k$ and $|\gamma_i| = 2^{q_i} p^{r_i}$ for a fixed odd prime p and for nonnegative integers r_i and q_i . For situations where this method is not applicable, Section 3 provides a simpler method of combining two grid graphs H_1 and H_2 where $rl(H)$ is merely equal to $rl(H_1)$ or $rl(H_2)$. Throughout, except for a few cases we have noted in the text, every result for finding Hamiltonian cycles can be easily adapted to find Hamiltonian paths as well.

2 Composition of cycles

Our primary method for finding Hamiltonian cycles with high run length consists of combining two or more Hamiltonian cycles to construct another Hamiltonian cycle in

their product; this method is modeled after those in [3] and [4]. Say the grid graph G_1 is the Cartesian product of cycles $\gamma_1 \times \gamma_2 \times \cdots \times \gamma_m$ and G_2 is $\gamma_{m+1} \times \gamma_{m+2} \times \cdots \times \gamma_k$, and let H_1 and H_2 be Hamiltonian cycles in G_1 and G_2 respectively. Then any Hamiltonian cycle H in $H_1 \times H_2$ is also a Hamiltonian cycle in $G = G_1 \times G_2$. We could consider H a cycle in the k -dimensional torus $\gamma_1 \times \gamma_2 \times \cdots \times \gamma_k$ or in the two-dimensional torus $H_1 \times H_2$, as demonstrated in Figure 1. If each γ_i in Figure 1 is a cycle on two vertices, and if H_1 is a cycle in $G_1 = \gamma_1 \times \gamma_2$ and H_2 is a cycle in $\gamma_3 \times \gamma_4 \times \gamma_5$, then an H which is Hamiltonian in $H_1 \times H_2$ is Hamiltonian in $G_1 \times G_2$. We can even determine the run length of H from this representation. Note that H takes three edges from the H_2 dimension followed by one from H_1 , and then repeats; if $rl(H_2) \geq 3$ and $rl(H_1) \geq 1$, then any adjacent four edges come from different dimensions. However, since $rl(H_2) = 2$, any three adjacent edges from H_2 may repeat a dimension, and so the run length may be no greater than two. We do know that any two adjacent edges are from different dimensions and thus that $rl(H)$ is at least 2. Note that, to avoid difficulty, we can construct such an H by assuming H_1 and H_2 are directed. In order to maximize $rl(H)$, we repeat an edge from the same dimension of H_i as seldom as possible.

This construction suggests the following theorem, couched in terms of $mrl(G)$, the maximum run length of any Hamiltonian cycle H in G . If $G = \gamma_1 \times \gamma_2 \times \cdots \times \gamma_k$, then $|G|$, the number of vertices in the grid graph G , is given by $\prod_{i=1}^k |\gamma_i|$.

Theorem 1. *Let G_1 and G_2 be grid graphs. If there exist s_1 and s_2 such that*

1. $\gcd(|G_1|, s_1) = 1$, $\gcd(|G_2|, s_2) = 1$, and
2. $\gcd(|G_1|, |G_2|) = s_1 + s_2$,

then $mrl(G_1 \times G_2) \geq mrl(G_1) + \left\lfloor \frac{s_2}{s_1} mrl(G_1) \right\rfloor$.

Proof. Choose Hamiltonian cycles H_1 and H_2 in G_1 and G_2 , respectively, with $rl(H_1) = mrl(G_1)$ and $rl(H_2) = mrl(G_2)$. Then we construct a Hamiltonian cycle H in $H_1 \times H_2$ with greatest possible run length. In [5], Trotter and Erdős show that a Hamiltonian cycle H exists in the directed Cartesian product $H_1 \times H_2$ exactly when there are integers s_1 and s_2 satisfying Conditions 1 and 2 above.

Curran and Witte explain in detail in [1] that Condition 2 ensures there is a cover of cycles where each point is contained in exactly one cycle exactly once. Any $s_1 + s_2$ consecutive edges contain s_1 from the dimension H_1 and s_2 from the dimension H_2 . In fact, any such subsequence of $s_1 + s_2$ consecutive edges uniquely determines the rest of the cycle or cover of cycles. Condition 1, on the other hand, ensures that H is a single Hamiltonian cycle rather than a cover of cycles since $\gcd(|G_1|, s_1) \cdot \gcd(|G_2|, s_2)$ gives the number of disjoint cycles in any such cover.

Subject to these conditions, we describe a construction to maximize $rl(H)$ based on the methods of Goddyn et al. [3]. First, we pick s_1 and s_2 so that s_1/s_2 is as close as possible to $rl(H_1)/rl(H_2)$. We then construct the cycle H by adding edges in steps so that, at the time of the j^{th} step, we have added x_{1j} from the direction of H_1 and x_{2j} from the direction of H_2 . At any point, the ratio x_{1j}/x_{2j} is as close as possible to s_1/s_2 without exceeding it. That is, if we start with an edge from the direction of H_2 , we then add as many as possible from the direction of H_1 , i.e., we add $\lfloor s_1/s_2 \rfloor$ edges, so $(x_{11}, x_{21}) = (\lfloor s_1/s_2 \rfloor, 1)$. Then we add another from H_2 followed by the appropriate

number from H_1 , so $(x_{12}, x_{22}) = (\lfloor 2s_1/s_2 \rfloor, 2)$, and so on. Ultimately, our first $s_1 + s_2$ edges will contain s_1 from H_1 and s_2 from H_2 . Any subpath of these $s_1 + s_2$ edges containing exactly $x_{1j} = rl(H)$ edges from H_1 will contain at least $x_{2i} \geq x_{1i}s_2/s_1 \geq \left\lfloor \frac{s_2}{s_1} rl(H_1) \right\rfloor$ edges from H_2 . That is,

$$rl(H) \geq rl(H_1) + \left\lfloor \frac{s_2}{s_1} rl(H_1) \right\rfloor.$$

Note that interchanging H_1 and H_2 may give a better bound. $\square$

Corollary 2. *If there exist s_1 and s_2 satisfying the conditions of Theorem 1, and if*

$$\frac{s_1}{s_2} = \frac{mrl(G_1)}{mrl(G_2)},$$

$$\text{then } mrl(G_1 \times G_2) \geq mrl(G_1) + mrl(G_2).$$

As an application of Theorem 1 and its Corollary 2, we have the following results.

Application 2.1. *If $G = \gamma_1 \times \gamma_2 \times \cdots \times \gamma_k$ is directed, then $mrl(G) < k$.*

Proof. Recall that the method introduced in Theorem 1 treats H_1 and H_2 as if they are directed. The theorem of Trotter and Erdős gives necessary and sufficient conditions for the existence of a Hamiltonian cycle in the product of exactly two directed cycles; thus, the conditions imposed are sufficient for the existence of cycles in G if it is undirected but are also necessary if it is considered as the product of directed cycles $H_1 \times H_2$. In our case, any cycle in G can be decomposed into a cycle in $H_1 \times H_2$ for some directed H_1 and H_2 . That is, if any directed graph $\gamma_1 \times \gamma_2 \times \cdots \times \gamma_i$ has maximum run length equal to i , then, for some ordering of the edges, there is a cycle in the graph which consists of the repeated subsequence of one edge from γ_1 , then one from γ_2 , $\dots$, one from γ_i , another one from γ_1 , etc. This implies first that $|G_1| = |\gamma_1||\gamma_2| \cdots |\gamma_{k-1}|$ is divisible by $i = k - 1$. It also means, however, that some cycle in the G given above can be decomposed into a cycle in $G_1 \times \gamma_k$ where $s_1 = mrl(G_1) = k - 1$. Then we know that $\gcd(|G_1|, k - 1) = 1$, which is a contradiction. $\square$

We may also make Corollary 2 more specific.

Corollary 3. *If s_1 and s_2 satisfy the conditions of Theorem 1 and if*

$$|s_2 \cdot rl(H_1) - s_1 \cdot rl(H_2)| \leq \max(s_1, s_2),$$

then

$$mrl(G_1 \times G_2) \geq mrl(G_1) + mrl(G_2) - 1.$$

Proof. Say $s_1 \geq s_2$. Then:

$$\left| \frac{s_2}{s_1} rl(H_1) - rl(H_2) \right| \leq 1$$

so

$$\left\lfloor \frac{s_2}{s_1} rl(H_1) \right\rfloor - rl(H_2) \leq 1.$$

The corollary follows. $\square$

Corollary 3 suggests another explicit application.

Application 2.2. Let p be a prime and let $G = \gamma_1 \times \gamma_2 \times \cdots \times \gamma_{3n}$ with $|G| = p^r$ for some positive integer r . Then $mrl(G) \geq n + 1$.

Proof. The base case was proved by Ruskey and Sawada [4], whose results imply $mrl(\gamma_1 \times \gamma_2 \times \gamma_3) \geq 2$ if $|\gamma_1||\gamma_2||\gamma_3| = p^r$.

For $n > 1$, consider $j = \lceil \frac{n}{2} \rceil$ and let $G_1 = \gamma_1 \times \gamma_2 \times \cdots \times \gamma_{3j}$ and $G_2 = \gamma_{3j+1} \times \gamma_{3j+2} \times \cdots \times \gamma_{3n}$ where $|G_1| = p^{r_1}$ and $|G_2| = p^{r_2}$ for p a prime and r_1, r_2 integers at least 3. Take Hamiltonian cycles H_1 in G_1 and H_2 in G_2 . Then $G = H_1 \times H_2$ and $\gcd(|H_1|, |H_2|) = p_0^{r_0}$ where $r_0 = \min(r_1, r_2)$. Assume inductively that $rl(H_1) = j + 1$ and $rl(H_2) = n - j + 1$. Then set

$$s_1 = \left\lfloor \frac{rl(H_1)}{rl(H_1) + rl(H_2)} \gcd(|H_1|, |H_2|) \right\rfloor + i = \left\lfloor \frac{j+1}{n+2} p^{r_0} \right\rfloor + i$$

with $i = 0$ or 1 so s_1 is not divisible by p . Also set $s_2 = p^{r_0} - s_1$ which is thus also not divisible by p . Then the conditions of Theorem 1 are satisfied, and

$$s_2 \cdot rl(H_1) - s_1 \cdot rl(H_2) = |s_2(j+1) - s_1(n-j+1)| \leq 2 \leq s_1,$$

so we may apply Corollary 3 to obtain $mrl(H_1 \times H_2) \geq n + 1$. $\square$

Application 2.3. Let p be a prime and $G = \gamma_1 \times \gamma_2 \times \cdots \times \gamma_{2n}$ where $|\gamma_i| = 2^{q_i} p^{r_i}$ for integers $q_i \geq 1$ and $r_i \geq 0$. Then $mrl(G) \geq n + 1$.

Proof. The method is very similar to that of Application 2.2. Again, Ruskey and Sawada's work [4] gives the base case $mrl(\gamma_1 \times \gamma_2) = 2$ for $|\gamma_1|$ and $|\gamma_2|$ even.

Set $j = \lceil \frac{n}{2} \rceil$ and take Hamiltonian cycles H_1 in $G_1 = \gamma_1 \times \gamma_2 \times \cdots \times \gamma_{2j}$ and H_2 in $G_2 = \gamma_{2j+1} \times \gamma_{2j+2} \times \cdots \times \gamma_{2n}$ so $G = H_1 \times H_2$ and $\gcd(|H_1|, |H_2|) = 2^{q_0} p^{r_0}$ for some integers $q_0 \geq 1$ and $r_0 \geq 0$. Assuming inductively that $rl(H_1) = j + 1$ and $rl(H_2) = n - j + 1$, we emulate Application 2.2 and set

$$s_1 = \left\lfloor \frac{j+1}{n+2} 2^{q_0} p^{r_0} \right\rfloor + i$$

for $i = 0, 1, 2$, or 3 chosen so s_1 not divisible by 2 or p . Then $s_2 = 2^{q_0} p^{r_0} - s_1$ is also not divisible by either. The conditions of Theorem 1 are met, and

$$|s_2(j+1) - s_1(n-j+1)| \leq 4 \leq s_1,$$

so Corollary 3 applies and $rl(H_1 \times H_2) \geq n + 1$. $\square$

3 Adding extra dimensions

We might ask, though, if there are easier ways of bounding $mrl(G)$. So far, we have primarily broken grid graphs into constituents $G_1 \times G_2$ where each constituent has maximum run length at least two, and where the run length of the whole is greater than that of either constituent. It is also possible, however, to ask if the Cartesian product of G and a k -dimensional cycle has maximum run length at least that of G . A method suggested by Ruskey and Sawada [4] is useful for one type of case.

Theorem 4. If $G = \gamma_1 \times \gamma_2 \times \cdots \times \gamma_k$, $|G|$ is even, and γ_{k+1} is a cycle, then $mrl(G \times \gamma_{k+1}) \geq mrl(G)$.

Proof. Without loss of generality, say $|\gamma_1| = 2n$. Then consider the embedding of H in G as a graph of connected lattice points $\mathbf{G}$ in $\mathbb{R}$, which is also a graph in $\mathbb{R}^k \times \{0\} \subset \mathbb{R}^{k+1}$. We wish to stack $|\gamma_{k+1}|$ copies of $\mathbf{G}$ on top of one another in $\mathbb{R}^{k+1}$ with every other one reversed. Reflect $\mathbf{G}$ exactly i times in the line $x_1 = n + 1/2$ to create $\mathbf{G}_i$ and then set it in $\mathbb{R} \times \{i\}$. Do this until there are $|\gamma_{k+1}|$ copies of $\mathbf{G}$ fixed immediately above each other with every other one reversed.




Figure 2: Two methods for extending cycles without decreasing run length are (a) extending the two-dimensional cycle G from $\gamma_1 \times \gamma_2$ into three dimensions where $|\gamma_3|$ is even and (b) extending the two-dimensional cycle G from $\gamma_1 \times \gamma_2$ to a three-dimensional one where $|\gamma_3|$ is a multiple of $|\gamma_2|$. In both cases, the solid lines represent the original cycle H , the dashed lines mark omitted edges, and the dotted lines mark added edges.

Pick any two edges $[v_{i,0}, v_{i,1}]$ and $[w_{i,0}, w_{i,1}]$ from the γ_1 direction located in the middle of $\mathbf{G}_0$. Then the point at $v_{i,0}$ is $\langle n, x_2, x_3, \dots, x_k, i \rangle$ and $v_{i,1} = v_{i,0} + \langle 1, 0, 0, \dots, 0 \rangle$, and, similarly, $w_{i,0} = \langle n, y_2, y_3, \dots, y_k, i \rangle$ and $w_{i,1} = w_{i,0} + \langle 1, 0, 0, \dots, 0 \rangle$. We assume that $x_i \neq y_i$. Note that these edges are preserved by the reflections, so $[v_{i,0}, v_{i,1}]$ is directly below $[v_{i+1,0}, v_{i+1,1}]$, and $[w_{i,0}, w_{i,1}]$ is also below $[w_{i+1,0}, w_{i+1,1}]$. Then remove these four edges and insert $[v_{i,0}, v_{i+1,0}]$, $[v_{i,1}, v_{i+1,1}]$, $[w_{i,0}, w_{i+1,0}]$, and $[w_{i,1}, w_{i+1,1}]$. This creates a cycle in $\bigcup \mathbf{G}_i$, which in turn induces a cycle H' in $G \times \gamma_{k+1}$. Where $rl(H) = mrl(G)$, any subsequence of $rl(H)$ adjacent edges in H with no more than one edge from any one dimension now corresponds to a sequence of $rl(H)$ adjacent edges in H' with still no more than one edge from any one direction. Thus, $mrl(G \times \gamma_{k+1}) \geq mrl(G)$ See Figure 2a. $\square$

As a continuation from Application 2.3, we have our next result.

Application 3.1. *Let $G = \gamma_1 \times \gamma_2 \times \cdots \times \gamma_k$ and $|\gamma_i| = 2^{q_i} p^{r_i}$, $q_i \geq 1, r_i \geq 0$. Then $mrl(G) \geq \lfloor k/2 \rfloor + 1$ for all k .*

Proof. If k is even, we may apply Corollary 3; if k is odd, $k - 1$ is even, and we may apply Theorem 4 and then Corollary 3 in succession. $\square$

A variation of this method is still available even if all the $|\gamma_i|$ are odd.

Theorem 5. *If $G = \gamma_1 \times \gamma_2 \times \cdots \times \gamma_k$ and $|\gamma_{k+1}|$ is a multiple of $|\gamma_j|$ for some $j \leq k$, then $mrl(G \times \gamma_{k+1}) \geq mrl(G)$.*

Proof. The proof is very similar to that of Theorem 5. The graph G embeds nicely as $\mathcal{G}_i$ in $\mathbb{R}^k \times \{i\}$; however, instead of reflecting $\mathcal{G}$, we translate it by i times the j^{th} unit vector. Then we remove some $[v_{1,0}, v_{1,1}]$ in $\mathcal{G}_1$ and all its images $[v_{i,0}, v_{i,1}]$ in $\mathcal{G}_i$. We attach $[v_{i,1}, v_{i+1,0}]$ to get a cycle in $\bigcup \mathcal{G}_i$ with the same run length as G . (Note that we must make sure that $v_{|\gamma_j|+1} = v_0$.) See Figure 2b. $\square$

Again, Theorem 5 allows us to generalize Application 2.2 as follows.

Application 3.2. *If $G = \gamma_1 \times \gamma_2 \times \cdots \times \gamma_k$ and $|G| = p^r$ for positive integer r , then $mrl(G) \geq \lfloor k/3 \rfloor + 1$ for all k .*

Proof. We already know that $mrl(G) \geq k/3 + 1$ for all k divisible by three. The rest follows by Theorem 5. $\square$

4 Conclusion and further work

This paper sets lower bounds on the maximum run length of some finite dimensional tori by giving methods for constructing Hamiltonian cycles of known run length. These methods produce Hamiltonian cycles in a torus $G = G_1 \times G_2$ from Hamiltonian cycles H in G_1 and G_2 .

The methods do not give very close upper and lower limits on $rl(H)$ and $mrl(G)$ and may be improved by future research. Additionally, there are many interesting cases not addressed in this paper, including: those cases where a path or a cover of disjoint cycles, but not a Hamiltonian cycle, can be found; grid graphs which are the cartesian products of paths or of paths and cycles; and grid graphs where some dimensions are directed and others are undirected. It would also be interesting to work out exactly which portion of the many possible grid graphs are covered by this work (since many of the results depend on the greatest common divisors of the sizes of the factor graphs). Additionally, the methods of Theorem 1 may be adjustable for any k -dimensional grid graph, perhaps using an extension of the idea of a diagonal (presented by Curran and Witte in [1]). Goddyn et al. [3] do present a version for a three-dimensional grid graph on eight vertices. The results in this paper also suggest a generalization of run length to j -variance, which we define to be the maximum r such that any sequence of r consecutive edges in a cycle H includes at most j edges from any one dimension (*N.B.*: 1-variance is the same as run length). Finally, a general but difficult challenge is finding methods for determining or estimating the maximum run length (or perhaps j -variance) of an arbitrary grid graph.

5 Acknowledgements

The work for this paper was mostly completed at the University of Minnesota, Duluth, funded by NSA grant MDA 904-02-1-0060 and NSF grant DMS-0137611. Many thanks are owed to the unflagging interest and input of Joe Gallian, Geir Helleloid, and Phil Matchett.

References

- [1] S. Curran, D. Witte. Hamiltonian paths in Cartesian products of directed cycles. *Cycles in Graphs* (Burnaby, B.C., 1982), 35-74, North-Holland Math.Stud., 115, North-Holland, Amsterdam, 1985.
- [2] L. Goddyn, P. Gvozdjak. Binary gray codes with long bit runs. *Electron. J. Combin.* 10 (2003), no.1, Research Paper 27, 12pp. (electronic).
- [3] L. Goddyn, G. Lawrence, E. Nemeth. Gray codes with optimized run lengths. *Utilitas Math.* 34 (1988), 179-192.
- [4] F. Ruskey, J. Sawada. Bent Hamilton cycles in d -dimensional grid graphs. *Electron. J. Combin.* 10 (2003), no. 1, Research Paper 1, 18pp. (electronic).
- [5] W. Trotter, P. Erdős. When the Cartesian product of directed cycles is Hamiltonian, *J. Graph Theory* 2 (1978), 137-142.